 <p>LEGACY HEALTH</p>	<p>Legacy Day Treatment Unit Provider's Orders</p> <p>Adult Ambulatory Infusion Order EFGARTIGIMOD ALFA (VYVGART AND VYVGART HYTRULO) FOR MYASTHENIA GRAVIS</p>	<p>Patient Name: _____</p> <p>Date of Birth: _____</p> <p>Med. Rec. No (TVC MRN Only): _____</p>
<p align="center">ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE</p>		

Anticipated Start Date: _____ **Patient to follow up with provider on date:** _____

*****This plan will expire after 365 days, unless otherwise specified below*****

Orders expire: _____

Weight: _____ kg **Height:** _____ cm

Allergies: _____

Diagnosis: _____

Diagnosis Code: _____ (please include primary and secondary diagnosis codes)

GUIDELINES FOR PRESCRIBING:


1. Send **FACE SHEET and H&P or most recent chart note.**
2. Patients should be up to date with all immunizations before initiating therapy. Avoid the use of live vaccines in patients during treatment.
3. Efgartigimod Alfa may increase the risk of infection. Delay treatment in patients with an active infection until the infection is resolved. Monitor for infection during treatment, and consider withholding treatment if infection develops.
4. Do NOT substitute efgartigimod alfa-hyaluronidase-qvfc (for SUBQ use) and efgartigimod alfa-fcab (for IV administration); products have different dosing and are NOT interchangeable

NURSING ORDERS (TREATMENT PARAMETERS):

1. TREATMENT PARAMETER – Hold infusion and contact provider if patient has signs or symptoms of infection.
2. Monitoring parameters depend on route selected in medications section:
 - a. If IV Infusion: Monitor patient for signs and symptoms of hypersensitivity reactions during infusion and for 1 hour following completion of infusion. Administer over 1 hour via 0.2 micron in-line filter.
 - b. If Subcutaneous injection: Monitor patient for signs and symptoms of hypersensitivity reactions during infusion and for 30 minutes following completion of injection. Administer using 12-inch tubing, PVC winged set. Choose an injection site on abdomen a minimum of 2 to 3 inches from the naval, avoiding areas with moles or scars, or where skin is red, bruised, or hard. Rotate injection sites for subsequent injections. Administer over a period of 30 to 90 seconds.

MEDICATIONS:

- Provider to Pharmacist Communication - Every visit, Administered once weekly x4 doses with subsequent cycles starting no sooner than 50 days from start of previous cycle (day 1, 8, 15, 22 every 50 days)
 - Efgartigimod alfa-fcab (VYVGART) 10 mg/kg (maximum dose: 1200 mg) in sodium chloride 0.9%, intravenous, over 1 hour, ONCE, weekly x 4 doses with subsequent cycles of once weekly x4 doses starting no sooner than 50 days from start of previous cycle.
 - Efgartigimod alfa 1008 mg/-hyaluronidase-qvfc 11,200 units (VYVGART HYTRULO) subcutaneous injection, ONCE, weekly x 4 doses with subsequent cycles of once weekly x4 doses starting no sooner than 50 days from start of previous cycle.

 <p>LEGACY HEALTH</p>	<p>Legacy Day Treatment Unit Provider's Orders</p> <p>Adult Ambulatory Infusion Order EFGARTIGIMOD ALFA (VYVGART AND VYVGART HYTRULO) FOR MYASTHENIA GRAVIS</p>	<p>Patient Name: _____</p> <p>Date of Birth: _____</p> <p>Med. Rec. No (TVC MRN Only): _____</p>
<p>ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE</p>		

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION - If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Refer to LH policy 906.6606 Initiation of Emergency Measures for Adult Oncology, Radiation Oncology and Infusion Clinic Patients
2. diphenhydramINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

Please check the appropriate box for the patient's preferred clinic location:

**Legacy Day Treatment Unit –
The Vancouver Clinic Building**
A department of Salmon Creek Medical Center
700 NE 87th Avenue, Suite 360
Vancouver, WA 98664
Phone number: 360-896-7070
Fax number: 360-487-5773

Legacy Emanuel Day Treatment Unit
A department of Emanuel Medical Center
501 N Graham Street, Suite 540
Portland, OR 97227
Phone number: 503-413-4608
Fax number: 503-413-4887

Legacy Salmon Creek Day Treatment Unit
Legacy Salmon Creek Medical Center
2121 NE 139th Street, Suite 110
Vancouver, WA 98686
Phone number: 360-487-1750
Fax number: 360-487-5773

Legacy STEPS Clinic
A department of Silverton Medical Center
Legacy Woodburn Health Center
1475 Mt Hood Ave
Woodburn, OR 97071
Phone number: 503-982-1280
Fax number: 503-225-8723

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Organization/Department: _____